1

Abstract

The algorithm checks the propositional formulas for patterns of unsatisfiability.

arXiv:cs/0703098v1 [cs.CC] 21 Mar 2007

Polynomial time algorithm for 3-SAT. Examples of use.

Sergey Gubin

February 1, 2008

1 Introduction

An $O(m^3)$-time algorithm for 3-SAT is described in [1]. This article illustrates that work with a few examples.

Let's summarize the algorithm. Let $c_1, c_2, \ldots, c_m$ be the clauses of a given 3-SAT instance. The algorithm detects whether there are compatible truth assignments among those truth assignments, which satisfy separate clauses. *Start.* Based on the clauses' truth-tables, let's build the clauses' compatibility matrices, grouped in the following box-matrix:

$$\begin{array}{cccc} c_1 : c_2 & c_1 : c_3 & \ldots & c_1 : c_m \\ & c_2 : c_3 & \ldots & c_2 : c_m \\ & & \ddots & \vdots \\ & & \ldots & c_{m-1} : c_m \end{array}$$

Step 1. Based on the first row of the box-matrix of compatibility matrices, let's deplete the compatibility matrices in the rows starting from the second. The resulting matrix is the following:

$$\begin{array}{cccc} c_1 : c_2 & c_1 : c_3 & \ldots & c_1 : c_m \\ & c_1 \wedge c_2 : c_1 \wedge c_3 & \ldots & c_1 \wedge c_2 : c_1 \wedge c_m \\ & & \ddots & \vdots \\ & & \ldots & c_1 \wedge c_{m-1} : c_1 \wedge c_m \end{array}$$

Next steps. Based on the second row of the resulting matrix, let's deplete the compatibility matrices in the rows starting from the third. And so on. After

*Author's email: sgubin@genesyslab.com

$m-2$ steps the matrix in the last row of the box-matrix of the compatibility matrices gets to be

$$c_1 \wedge c_2 \wedge \ldots \wedge c_{m-1} : c_1 \wedge c_2 \wedge \ldots \wedge c_m$$

Thus, the given clauses are compatible (the given 3-SAT instance is satisfiable) iff the algorithm does not create matrices filled with $false$ completely.

In the tables below, "1" is used for $true$ and empty string for $false$. Some extra white space in the article is due to allocation of the tables.

2 Example 1

The 3-SAT instance from [2]:

$$f = (p \vee q \vee r) \wedge (\bar{p} \vee q \vee \bar{r}) \wedge (p \vee \bar{q} \vee s) \wedge (\bar{p} \vee \bar{r} \vee \bar{s}).$$

Let's use the algorithm directly as it was described in Theorem 1 in [1]. Clauses' truth-tables:

$c_1 =$

#	p	q	r	$p \vee q \vee r$
1				
2			1	1
3		1		1
4		1	1	1
5	1			1
6	1		1	1
7	1	1		1
8	1	1	1	1

$c_2 =$

#	p	q	r	$\bar{p} \vee q \vee \bar{r}$
1				1
2			1	1
3		1		1
4		1	1	1
5	1			1
6	1		1	
7	1	1		1
8	1	1	1	1

$c_3 =$

#	p	q	s	$p \vee \bar{q} \vee s$
1				1
2			1	1
3		1		
4		1	1	1
5	1			1
6	1		1	1
7	1	1		1
8	1	1	1	1

$c_4 =$

#	p	r	s	$\bar{p} \vee \bar{r} \vee \bar{s}$
1				1
2			1	1
3		1		1
4		1	1	1
5	1			1
6	1		1	1
7	1	1		1
8	1	1	1	

Start. Based on the truth-tables, we calculate clauses' compatibility matrices. Compatibility matrix $c_i : c_j$ shows which meanings of clauses c_i and c_j are

compatible, $i < j;\ i = 1, 2, 3;\ j = 2, 3, 4$:

$c_1 : c_2$	$c_1 : c_3$	$c_1 : c_4$
	$c_2 : c_3$	$c_2 : c_4$
		$c_3 : c_4$

The matrices are shown below.
Step 1. Let's deplete compatibility matrices $c_i : c_j,\ i \geq 2$. These matrices are allocated in the second and the third rows on the pictures below. Let's get rid of those elements $(\alpha\beta)$ of these matrices, which are not compatible with any truth assignment for clause c_1:

$c_1 : c_2$	$c_1 : c_3$	$c_1 : c_4$
	$c_1 \wedge c_2 : c_1 \wedge c_3$	$c_1 \wedge c_2 : c_1 \wedge c_4$
		$c_1 \wedge c_3 : c_1 \wedge c_4$

For example, element (1,1) of compatibility matrix $c_3 : c_4$ is associated with the first meaning of clause c_3 and the first meaning of clause c_4. The numbers are taken in accordance with truth-tables for the clauses. But there is not any meaning of clause c_1, which would be compatible simultaneously with the first meaning of clause c_3 and the first meaning of clause c_4. Thus, element (1,1) of compatibility matrix $c_3 : c_4$ must be eliminated.

The resulting matrices are shown below. Because there is not any $false$-matrix, let's continue.
Step 2. Let's deplete compatibility matrix $c_3 : c_4$. The matrix is allocated in the third row. Let's get rid of those elements $(\alpha\beta)$, for which there is not any truth assignment for clause c_2 left, which would be simultaneously compatible with α-th truth assignment for c_3 and with β-th truth assignment for clause $c_{4.}$. Let's mention that, after step 1, matrix $c_3 : c_4$ contains only those couples, which are compatible with clause c_1:

$c_1 : c_2$	$c_1 : c_3$	$c_1 : c_4$
	$c_1 \wedge c_2 : c_1 \wedge c_3$	$c_1 \wedge c_2 : c_1 \wedge c_4$
		$c_1 \wedge c_2 \wedge c_3 : c_1 \wedge c_2 \wedge c_4$

The resulting matrix is shown below. Because it is not a $false$-matrix, the example is a satisfiable 3-SAT instance.

Start. Clauses' compatibility matrices:

c_1 / c_2

	1	2	3	4	5	6	7	8
1								
2		1						
3			1					
4				1				
5					1			
6								
7							1	
8								1

c_1 / c_3

	1	2	3	4	5	6	7	8
1								
2	1	1						
3				1				
4				1				
5					1	1		
6					1	1		
7							1	1
8							1	1

c_1 / c_4

	1	2	3	4	5	6	7	8
1								
2			1	1				
3	1	1						
4			1	1				
5					1	1		
6							1	
7					1	1		
8							1	

c_2 / c_3

	1	2	3	4	5	6	7	8
1	1	1						
2	1	1						
3				1				
4				1				
5					1	1		
6								
7							1	1
8							1	1

c_2 / c_4

	1	2	3	4	5	6	7	8
1	1	1						
2			1	1				
3	1	1						
4			1	1				
5					1	1		
6								
7					1	1		
8							1	

c_3 / c_4

	1	2	3	4	5	6	7	8
1	1		1					
2		1		1				
3								
4		1		1				
5					1		1	
6						1		
7					1		1	
8						1		

Step 1. Compatibility matrices in the second and the third rows are depleted by clause c_1:

c_1

	c_2							
	1	2	3	4	5	6	7	8
1								
2		1						
3			1					
4				1				
5					1			
6								
7							1	
8								1

	c_3							
	1	2	3	4	5	6	7	8
1								
2	1	1						
3				1				
4				1				
5					1	1		
6					1	1		
7							1	1
8							1	1

	c_4							
	1	2	3	4	5	6	7	8
1								
2			1	1				
3	1	1						
4			1	1				
5					1	1		
6							1	
7					1	1		
8							1	

c_2

	c_3							
	1	2	3	4	5	6	7	8
1								
2	1	1						
3				1				
4				1				
5					1	1		
6								
7							1	1
8							1	1

	c_4							
	1	2	3	4	5	6	7	8
1								
2			1	1				
3	1	1						
4			1	1				
5					1	1		
6								
7					1	1		
8							1	

c_3

	c_4							
	1	2	3	4	5	6	7	8
1			1					
2				1				
3								
4		1		1				
5					1		1	
6						1		
7					1		1	
8						1		

Step 2. Compatibility matrix $c_3 : c_4$ in the third row is depleted by clauses c_1 and c_2:

c_1 \ c_2

	1	2	3	4	5	6	7	8
1								
2		1						
3			1					
4				1				
5					1			
6								
7							1	
8								1

c_3

	1	2	3	4	5	6	7	8
1								
2	1	1						
3				1				
4				1				
5					1	1		
6					1	1		
7							1	1
8							1	1

c_4

	1	2	3	4	5	6	7	8
1								
2			1	1				
3	1	1						
4			1	1				
5					1	1		
6							1	
7					1	1		
8							1	

c_3 \ c_4

	1	2	3	4	5	6	7	8
1								
2			1	1				
3	1	1						
4			1	1				
5					1	1		
6								
7					1	1		
8							1	

c_2 \ c_4

	1	2	3	4	5	6	7	8
1			1					
2				1				
3								
4		1		1				
5					1			
6						1		
7					1		1	
8						1		

Because there is not any $false$-matrix, the 3-SAT instance is satisfiable.

All solutions. In the same way, let's move backward and get rid of all elements in the box-matrix of the compatibility matrices except those satisfying the given formula:

c_1

	c_2							
	1	2	3	4	5	6	7	8
1								
2		1						
3			1					
4				1				
5					1			
6								
7							1	
8								1

	c_3							
	1	2	3	4	5	6	7	8
1								
2	1	1						
3				1				
4				1				
5					1	1		
6								
7							1	1
8							1	

	c_4							
	1	2	3	4	5	6	7	8
1								
2			1	1				
3		1						
4				1				
5					1	1		
6								
7					1	1		
8							1	

c_2

	c_3							
	1	2	3	4	5	6	7	8
1								
2	1	1						
3				1				
4				1				
5					1	1		
6								
7							1	1
8							1	

	c_4							
	1	2	3	4	5	6	7	8
1								
2			1	1				
3		1						
4				1				
5					1	1		
6								
7					1	1		
8							1	

c_3

	c_4							
	1	2	3	4	5	6	7	8
1			1					
2				1				
3								
4		1		1				
5					1			
6						1		
7					1		1	
8						1		

Brute force method. Let's compare the solution with the solution obtained with the brute force method. The brute force solution is shown in the table below.

#	p	q	r	s	c_1	c_2	c_3	c_4	f	$c_1 : c_2$	$c_1 : c_3$	$c_1 : c_4$	$c_2 : c_3$	$c_2 : c_4$	$c_3 : c_4$
1						1	1	1							
2				1		1	1	1							
3			1		1	1	1	1	1	2 : 2	2 : 1	2 : 3	2 : 1	2 : 3	1 : 3
4			1	1	1	1	1	1	1	2 : 2	2 : 2	2 : 4	2 : 2	2 : 4	2 : 4
5		1			1	1		1							
6		1		1	1	1	1	1	1	3 : 3	3 : 4	3 : 2	3 : 4	3 : 2	4 : 2
7		1	1		1	1		1							
8		1	1	1	1	1	1	1	1	4 : 4	4 : 4	4 : 4	4 : 4	4 : 4	4 : 4
9	1				1	1	1	1	1	5 : 5	5 : 5	5 : 5	5 : 5	5 : 5	5 : 5
10	1			1	1	1	1	1	1	5 : 5	5 : 6	5 : 6	5 : 6	5 : 6	6 : 6
11	1		1		1	1		1							
12	1		1	1		1									
13	1	1			1	1	1	1	1	7 : 7	7 : 7	7 : 5	7 : 7	7 : 5	7 : 5
14	1	1		1	1	1	1	1	1	7 : 7	7 : 8	7 : 6	7 : 8	7 : 6	8 : 6
15	1	1	1		1	1	1	1	1	8 : 8	8 : 7	8 : 7	8 : 7	8 : 7	7 : 7
16	1	1	1	1	1	1	1								

3 Example 2

The 3-SAT instance from [3]:

$$f = (p \vee q \vee r) \wedge (p \vee q \vee \bar{r}) \wedge (\bar{p} \vee s) \wedge (\bar{p} \vee \bar{s}) \wedge \bar{q}.$$

Let's use the formula (4) from [1] to iterate the compatibility matrices.

For example, compatibility matrices $C_{14,0} = c_1 : c_4$, $C_{15,0} = c_1 : c_5$, and $C_{45,0} = c_4 : c_5$ are (see below)

$C_{14,0} =$

1	1		
1	1		
1	1		
		1	
		1	
		1	
		1	

$C_{15,0} =$

1	
1	
1	

$C_{45,0} =$

1	
1	
1	

Then, formula (4) from [1] works in the following way:

$$C_{45,1} = (C_{14,0}^T \times C_{15,0}) \wedge C_{45,0} =$$

$$= (\begin{array}{|c|c|c|c|c|c|c|c|}\hline & 1 & 1 & 1 & & & & \\\hline & 1 & 1 & 1 & & & & \\\hline & & & & 1 & 1 & 1 & 1 \\\hline & & & & & & & \\\hline\end{array} \times \begin{array}{|c|c|}\hline & \\\hline 1 & \\\hline & \\\hline & \\\hline 1 & \\\hline 1 & \\\hline & \\\hline & \\\hline\end{array}) \wedge \begin{array}{|c|c|}\hline 1 & \\\hline 1 & \\\hline 1 & \\\hline & \\\hline\end{array} =$$

$$= \begin{array}{|c|c|}\hline 1 & \\\hline 1 & \\\hline 1 & \\\hline & \\\hline\end{array} \wedge \begin{array}{|c|c|}\hline 1 & \\\hline 1 & \\\hline 1 & \\\hline & \\\hline\end{array} = \begin{array}{|c|c|}\hline 1 & \\\hline 1 & \\\hline 1 & \\\hline & \\\hline\end{array}.$$

Multiplication of Boolean matrices can be defined in different ways. The rules used here are the rules of numeric matrices' multiplication but with replacement of numeric multiplications and sums with conjunctions and disjunctions appropriately [1].

Let's return to the problem. Clauses' truth-tables:

$c_1 =$

#	p	q	r	$p \vee q \vee r$
1				
2			1	1
3		1		1
4		1	1	1
5	1			1
6	1		1	1
7	1	1		1
8	1	1	1	1

$c_2 =$

#	p	q	r	$p \vee q \vee \bar{r}$
1				1
2			1	
3		1		1
4		1	1	1
5	1			1
6	1		1	1
7	1	1		1
8	1	1	1	1

$c_3 =$

#	p	s	$\bar{p} \vee s$
1			1
2		1	1
3	1		
4	1	1	1

$c_4 =$

#	p	s	$\bar{p} \vee \bar{s}$
1			1
2		1	1
3	1		1
4	1	1	

$c_5 =$

#	q	$\bar{q}$
1		1
2	1	

Start. Compatibility matrices:

c_1

	c_2							
	1	2	3	4	5	6	7	8
1								
2								
3			1					
4				1				
5					1			
6						1		
7							1	
8								1

c_3			
1	2	3	4
1	1		
1	1		
1	1		
			1
			1
			1
			1

c_4			
1	2	3	4
1	1		
1	1		
1	1		
		1	
		1	
		1	
		1	

c_5	
1	2
1	
1	
1	

c_2

	c_3			
	1	2	3	4
1	1	1		
2				
3	1	1		
4	1	1		
5				1
6				1
7				1
8				1

c_4			
1	2	3	4
1	1		
1	1		
1	1		
		1	
		1	
		1	
		1	

c_5	
1	2
1	
1	
1	

c_3

	c_4			
	1	2	3	4
1	1			
2		1		
3				
4				

c_5	
1	2
1	
1	
1	

c_4

	c_5	
	1	2
1	1	
2	1	
3	1	
4		

Step 1. Compatibility matrices depleted by clause c_1 with formula (4) from [1]:

c_1

	c_2							
	1	2	3	4	5	6	7	8
1								
2								
3			1					
4				1				
5					1			
6						1		
7							1	
8								1

	c_3			
	1	2	3	4
1				
2	1	1		
3	1	1		
4	1	1		
5				1
6				1
7				1
8				1

	c_4			
	1	2	3	4
1				
2	1	1		
3	1	1		
4	1	1		
5			1	
6			1	
7			1	
8			1	

	c_5	
	1	2
1		
2	1	
3		
4		
5	1	
6	1	
7		
8		

c_2

	c_3			
	1	2	3	4
1				
2				
3	1	1		
4	1	1		
5				1
6				1
7				1
8				1

	c_4			
	1	2	3	4
1				
2				
3	1	1		
4	1	1		
5			1	
6			1	
7			1	
8			1	

	c_5	
	1	2
1		
2		
3		
4		
5	1	
6	1	
7		
8		

c_3

	c_4			
	1	2	3	4
1	1			
2		1		
3				
4				

	c_5	
	1	2
1	1	
2	1	
3		
4	1	

c_4

	c_5	
	1	2
1	1	
2	1	
3	1	
4		

Step 2. Compatibility matrices depleted by clause c_2 with formula (4) from [1]:

c_1	c_2								c_3				c_4				c_5	
		1	2	3	4	5	6	7	8	1	2	3	4	1	2	3	4	1	2
	1																		
	2									1	1			1	1			1	
	3			1						1	1			1	1				
	4				1					1	1			1	1				
	5					1							1			1		1	
	6						1						1			1		1	
	7							1					1			1			
	8								1				1			1			

c_2	c_3				c_4				c_5	
	1	2	3	4	1	2	3	4	1	2
1										
2										
3	1	1			1	1				
4	1	1			1	1				
5				1			1		1	
6				1			1		1	
7				1			1			
8				1			1			

c_3	c_4				c_5	
	1	2	3	4	1	2
1	1					
2		1				
3						
4					1	

c_4	c_5	
	1	2
1		
2		
3	1	
4		

Step 3. Compatibility matrices depleted by clause c_3 with formula (4) from [1]:

c_1

	c_2								c_3				c_4				c_5	
	1	2	3	4	5	6	7	8	1	2	3	4	1	2	3	4	1	2
1																		
2									1	1			1	1			1	
3			1						1	1			1	1				
4				1					1	1			1	1				
5					1							1			1		1	
6						1						1			1		1	
7							1					1			1			
8								1				1			1			

c_2

	c_3				c_4				c_5	
	1	2	3	4	1	2	3	4	1	2
1										
2										
3	1	1			1	1				
4	1	1			1	1				
5				1			1		1	
6				1			1		1	
7				1			1			
8				1			1			

c_3

	c_4				c_5	
	1	2	3	4	1	2
1	1					
2		1				
3						
4					1	

c_4

	c_5	
	1	2
1		
2		
3		
4		

The compatibility matrix $C_{45,3} = c_4 : c_5$ is a $false$-matrix. Thus, the given 3-SAT instance is unsatisfiable.

Let's emphasize that initial compatibility matrix $c_4 : c_5$ was depleted by clauses c_4 and c_5. It contained (its $true$-elements were associated with) only compatible true assignments of clauses c_4 and c_5. The algorithm's steps 1, 2, and 3 cleaned those true assignments to make them compatible with clauses c_1, c_2, and c_3, also. It so happened that there were no possibilities left after that cleaning in this example.

The tree of possibilities collapses due to formula (4) [1]. The Boolean matrices' multiplication "totals" the possibilities. And the Boolean matrices' conjunction "clears" the totals. Sure, some totals can be empty, too.

For example, let's take $(3, 1)$-element of matrix $C_{45,3}$ (Step 3). The multiplication part of formula (4) [1] totals all possibilities sorted over the third columns of matrix $C_{34,2}$ and the first column of matrix $C_{35,2}$ into only one possibility:

$$false \wedge false \vee false \wedge false \vee false \wedge false \vee false \wedge true = false.$$

That means that there are no possibilities left. Then, the conjunction part of the formula (4) [1] clears the total possibility against possibilities already accounted for in the $(3, 1)$-element of matrix $C_{45,2}$:

$$false \wedge true = false.$$

4 Example 3

Let's see how the algorithm deals with 2-SAT. The 2-SAT instance from [4]:

$$f = \bar{p} \wedge \bar{q} \wedge \bar{r} \wedge (p \vee q) \wedge (p \vee r) \wedge (q \vee r).$$

Let's use formula (4) from [1] to iterate the compatibility matrices.

Clauses' truth-tables:

$c_1 =$

#	p	$\bar{p}$
1		1
2	1	

$c_2 =$

#	q	$\bar{q}$
1		1
2	1	

$c_3 =$

#	r	$\bar{r}$
1		1
2	1	

$c_4 =$

#	p	q	$p \vee q$
1			
2		1	1
3	1		1
4	1	1	1

$c_5 =$

#	p	r	$p \vee r$
1			
2		1	1
3	1		1
4	1	1	1

$c_6 =$

#	q	r	$q \vee r$
1			
2		1	1
3	1		1
4	1	1	1

Start. Compatibility matrices:

c_1

	c_2		c_3		c_4				c_5				c_6			
		1	2	1	2	1	2	3	4	1	2	3	4	1	2	3	4
1	1		1			1					1				1	1	1
2																	

c_2

	c_3		c_4				c_5				c_6			
		1	2	1	2	3	4	1	2	3	4	1	2	3	4
1	1					1			1	1	1		1		
2															

c_3

	c_4					c_5				c_6			
		1	2	3	4	1	2	3	4	1	2	3	4
1	1		1	1	1			1				1	
2													


c_4

	c_5					c_6			
		1	2	3	4	1	2	3	4
1	1								
2			1					1	1
3				1	1		1		
4				1	1			1	1

c_5

	c_6				
		1	2	3	4
1	1				
2			1		1
3				1	
4			1		1

Step 1. Compatibility matrices depleted by clause c_1 with formula (4) from [1]:



Here, patterns of unsatisfiability arose - matrices $c_2 : c_4$ and $c_3 : c_5$ are $false$-matrices. Thus, the 2-SAT instance is unsatisfiable.

Let's mention that $false$-matrix $c_2 : c_4$ in Step 1 (matrix $C_{24,1}$) shows that formula f contains an unsatisfiable part

$$c_1 \wedge c_2 \wedge c_4 = \bar{p} \wedge \bar{q} \wedge (p \vee q).$$

And $false$-matrix $c_3 : c_5$ in the Step 1 (matrix $C_{35,1}$) shows that formula f contains an unsatisfiable part

$$c_1 \wedge c_3 \wedge c_5 = \bar{p} \wedge \bar{r} \wedge (p \vee r).$$

Formula f contains one more unsatisfiable part

$$c_2 \wedge c_3 \wedge c_6 = \bar{q} \wedge \bar{r} \wedge (q \vee r).$$

But that would be detected only in Step 2 after the depletion of compatibility matrix $c_3 : c_6$ by clause c_2.

The phenomena noticed here are not accidental.

Theorem 1. *(Structure of unsatisfiable SAT instances)*
1). If the algorithm stops after k steps, then the given propositional formula contains $k + 2$ clauses that create an unsatisfiable formula.
2). If the given propositional formula contains k clauses that create an unsatisfiable formula, then there is such an enumeration of clauses that the algorithm will stop in $k - 2$ steps.

For example, the algorithm will not even iterate for the formula

$$x \wedge \bar{x}.$$

It will detect that the formula is unsatisfiable on its initial step "Start".

5 Example 4

The following 3-SAT instances:

$$f_1 = (p \vee q \vee r) \wedge (p \vee q \vee \bar{r}) \wedge (p \vee \bar{q} \vee r) \wedge (p \vee \bar{q} \vee \bar{r}) \wedge (\bar{p} \vee q \vee r) \wedge (\bar{p} \vee q \vee \bar{r}) \wedge (\bar{p} \vee \bar{q} \vee r),$$

$$f_2 = f_1 \wedge (\bar{p} \vee \bar{q} \vee \bar{r}).$$

Clauses and their truth-tables:

	c_i
c_1	$p \vee q \vee r$
c_2	$p \vee q \vee \bar{r}$
c_3	$p \vee \bar{q} \vee r$
c_4	$p \vee \bar{q} \vee \bar{r}$
c_5	$\bar{p} \vee q \vee r$
c_6	$\bar{p} \vee q \vee \bar{r}$
c_7	$\bar{p} \vee \bar{q} \vee r$
c_8	$\bar{p} \vee \bar{q} \vee \bar{r}$

#	p	q	r	c_1	c_2	c_3	c_4	c_5	c_6	c_7	c_8
1					1	1	1	1	1	1	1
2			1	1		1	1	1	1	1	1
3		1		1	1		1	1	1	1	1
4		1	1	1	1	1		1	1	1	1
5	1			1	1	1	1		1	1	1
6	1		1	1	1	1	1	1		1	1
7	1	1		1	1	1	1	1	1		1
8	1	1	1	1	1	1	1	1	1	1	

Due to the selected enumeration of the strings in the truth-tables, the compatibility matrices are the following diagonal Boolean matrices:

$$C_{ij} = (c_i \wedge c_j) = diag(b_{ij1}, b_{ij2}, b_{ij3}, b_{ij4}, b_{ij5}, b_{ij6}, b_7, b_{ij8}),$$

- where $i = 1, 2, \ldots, 7;\ j = 2, 3, \ldots, 8$; and

$$b_{ijk} = \begin{cases} false & k = i \vee k = j \\ true & k \neq i \wedge k \neq j \end{cases},$$

- where $k = 1, 2, \ldots, 8$. Thus, each iteration of the algorithm will eliminate (make it $false$) from the initial compatibility matrices at least one diagonal element. After $n \leq 8 - 2 = 6$ iterations, the result will contain only those elements, which pertain their value $true$. Those elements are common for all compatibility matrices. For formula f_1, such element is

$$p = q = r = true.$$

But for formula f_2, there are no such elements. Thus, f_1 is satisfiable, and f_2 is not.

References

[1] Sergey Gubin. A Polynomial Time Algorithm for 3-SAT. http://www.arxiv.org/PS_cache/cs/pdf/0701/0701023.pdf

[2] Stephen Cook. The P versus NP problem. http://www.claymath.org/millennium/P_vs_NP/pvsnp.pdf

[3] Craig Feinstein. Email to author.

[4] Blake Hegerle. A Counterexample to a Proposed Proof of P = NP by S. Gubin. http://arxiv.org/PS_cache/cs/pdf/0701/0701033.pdf